Table 12C 2001 Pediatric Nutrition Surveillance (1) Kentucky

Page: 1 Run Date: 8/20/2002

Summary of Trends in Growth and Anemia Indicators Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)													
	Bi	rthweight ((4)		Anemia Low Hb/Hct								
Year	Low High				Short Under- Over- Stature weight weight (5) (6) (6) (6) (6) ≤ 2 Yrs Overweight (6)						(7)		
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%	
2001	31,762	9.5	7.9	113,838	5.0	5.7	14.2	48,713	15.7	14.4	77,701	12.3	
2000	33,613	9.5	8.5	112,873	4.6	6.3	13.5	47,657	15.4	13.9	76,598	11.2	
1999	11,780	9.3	8.3	39,130	4.5	6.4	13.3	17,851	16.1	13.5	24,723	10.5	
1998	23,911	9.1	8.5	100,855	4.7	6.0	12.7	48,075	15.1	12.1	68,824	11.2	
1997	26,884	8.9	8.7	111,523	4.8	6.2	12.0	52,606	14.7	11.5	73,732	13.0	
1996	32,550	9.0	9.1	122,881	4.9	6.3	11.6	55,008	14.5	11.0	85,062	13.1	
1995	31,197	8.7	8.8	120,246	5.4	6.4	11.1	53,344	14.4	10.2	91,604	14.4	
1994	32,318	8.7	9.2	118,238	5.6	6.4	11.2	50,388	14.3	10.1	89,046	14.8	
1993	26,816	8.7	9.1	98,920	5.6	6.7	10.6	43,531	13.4	9.4	75,879	14.8	
1992	9,244	8.2	9.2	39,037	5.9	6.5	9.9	18,006	12.9	8.5	31,027	13.6	

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and height-for-age for age and height-for-age for children 2 years of age and height-for-age for age age and heig

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C 2001 Pediatric Nutrition Surveillance (1) Kentucky

Page: 2 Run Date: 8/20/2002

Summary of Trends in Growth and Anemia Indicators Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)													
	Bi	irthweight ((4)	Height and Weight								Anemia Low Hb/Hct	
Year	Low High				Short Under- Over- Stature weight weight (5) (6) (6) (6) (6) ≤ 2 Yrs Overweight (6)						(7)		
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number 8	% 5th-<95th	% ≥95th	Number	%	
1991	24,422	8.5	9.3	87,953	6.2	6.7	9.7	37,114	12.5	8.0	66,344	14.2	
1990	27,863	8.4	9.3	97,214	6.1	6.9	9.9	39,198	12.4	8.4	69,312	12.7	
1989	21,053	8.1	9.6	76,180	6.1	7.0	9.3	31,985	12.6	7.9	56,436	11.9	
1988	25,911	8.4	9.5	78,950	7.5	7.5	9.8	30,733	12.2	8.0	57,116	10.5	
1987	24,503	8.9	8.8	70,534	9.0	7.5	10.6	27,085	11.3	7.6	52,126	11.7	
1986	23,303	9.0	8.9	71,888	8.8	7.8	10.3	27,049	10.9	7.6	53,401	13.0	
1985	22,495	9.1	8.8	73,473	8.6	6.9	10.5	28,457	11.2	7.4	59,738	15.2	
1984	21,168	9.3	8.9	67,645	9.0	7.1	10.2	25,916	11.8	7.0	55,323	18.5	
1983	20,357	9.5	8.5	61,917	9.2	8.0	9.6	22,203	11.3	6.6	49,822	23.3	
1982	14,165	9.7	9.4	56,632	8.4	7.3	8.9	19,699	11.5	7.4	44,190	20.2	

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 3 years of age and height-for-age

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C 2001 Pediatric Nutrition Surveillance (1) Kentucky

Page: 3 Run Date: 8/20/2002

Summary of Trends in Growth and Anemia Indicators Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)													
	Bi	rthweight (4)	Height and Weight								Anemia Low Hb/Hct	
Year		Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			(7)		
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>></u> 95th	Number 8	% 5th-<95th	% <u>></u> 95th	Number	%	
1981	16,002	9.4	9.3	61,302	8.4	7.1	9.2	21,373	11.8	7.6	47,553	20.9	
1980	16,486	9.2	9.3	64,223	8.6	7.7	9.7	23,476	11.9	7.8	50,141	21.8	
1979	12,570	9.3	9.2	46,041	8.8	7.0	10.7	17,027	13.0	8.2	35,044	23.2	
1978	10,384	9.0	9.3	41,525	8.7	7.5	10.4	16,388	12.7	7.9	31,986	23.8	
1977	9,009	10.2	8.9	34,075	10.3	8.6	10.8	13,516	12.3	8.0	25,881	26.2	
1976	4,564	10.9	7.8	21,168	10.7	7.1	12.9	8,958	13.0	9.7	16,664	17.7	
1975	3,950	10.7	7.8	18,222	11.1	8.3	12.5	6,799	12.8	10.0	14,429	20.6	
1974	2,868	11.4	9.4	14,230	13.2	8.6	13.9	5,595	14.8	10.6	11,503	27.3	
1973	189	9.5	11.6	586	8.4	8.0	13.1	250	11.2	7.2	487	15.4	



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and height-for-age for age and height-for-age for children 2 years of age and height-for-age for age age and heig

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.